

University of Groningen

Multinuclear non-heme iron complexes for double-strand DNA cleavage

Megens, Rik P.; van den Berg, Tieme A.; de Bruijn, A. Dowine; Feringa, Ben L.; Roelfes, Gerard

Published in:
 Chemistry

DOI:
[10.1002/chem.200801409](https://doi.org/10.1002/chem.200801409)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Megens, R. P., van den Berg, T. A., de Bruijn, A. D., Feringa, B. L., & Roelfes, G. (2009). Multinuclear non-heme iron complexes for double-strand DNA cleavage. *Chemistry*, 15(7), 1723-1733.
<https://doi.org/10.1002/chem.200801409>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

CHEMISTRY

A EUROPEAN JOURNAL

Supporting Information

© Copyright Wiley-VCH Verlag GmbH & Co. KGaA, 69451 Weinheim, 2008

Multinuclear Non-Heme Iron Complexes for Double Strand DNA Cleavage

Rik P. Megens, Tieme A. van den Berg, A. Dowine de Bruijn, Ben L. Feringa* and Gerard Roelfes*^[a]

*Stratingh Institute for Chemistry
University of Groningen
Nijenborgh 4, 9747 AG Groningen (The Netherlands)*

Rik P. Megens, Tieme A. van den Berg, A. Dowine de Bruijn, Ben L. Feringa* and Gerard Roelfes*

Stratingh Institute for Chemistry
University of Groningen
Nijenborgh 4, 9747 AG Groningen (The Netherlands)
Fax: (+31) 50-363-4296
E-mail: j.g.roelfes@rug.nl; b.l.feringa@rug.nl

Instrumentation

NMR spectra were recorded on a Varian Mercury Plus 200 (^1H NMR at 200 MHz, ^{13}C NMR at 50.3 MHz), a Varian VXR-300 (^1H NMR at 300 MHz, ^{13}C NMR at 75.5 MHz), or on a Varian Mercury Plus 400 (^1H NMR at 400 MHz, ^{13}C NMR at 100 MHz). Chemical shifts (δ) are denoted in ppm and referenced to the residual solvent peak unless stated otherwise (CDCl_3 , ^1H δ = 7.24, ^{13}C δ = 77.0; CD_3OD , ^1H δ = 3.30, ^{13}C δ = 49.0; CD_3CN , ^1H δ = 1.93, D_2O ^1H δ = 4.79). The splitting patterns are designated as follows: s (singlet), d (doublet), dd (double doublet), t (triplet), q (quartet), m (multiplet), and br (broad). Coupling constants (J) between two nuclei separated by n chemical bonds are denoted in hertz (Hz).

Chemical ionisation mass spectra (MS-Cl^+), electron impact (MS-EI^+), and exact mass determination (HRMS) were recorded on a AEI MS-902. Electrospray ionisation mass spectrometry (MS-ESI^+) was performed on a Triple Quadrupole LC-MS-MS mass spectrometer (API 3000, Perkin-Elmer Sciex Instruments).

Elemental analysis was performed on an EuroVector CHNS-O Elemental Analyzer Euro EA 3000. Melting points were recorded on a Büchi B-545 melting point apparatus. UV-Vis spectra were recorded on a Hewlett-Packard 8453 diode array or Jasco V-570 spectrophotometer. High resolution ESI-MS was performed on a Applied Biosystems Q-STAR mass spectrometer.

HPLC separations

Reversed-phase high-performance liquid chromatography (RP-HPLC) was performed on a Shimadzu LC-10AD vp at 35°C, which was equipped with a photodiode array detector (SPD M10A) and a fraction collector (Shimadzu FRC 10A). A Waters 2690 Separations Module equipped with a photodiode array detector (Waters 996) and a Micromass ZMD Quadrupole mass spectrometer at room temperature was used for LC-MS analysis.

Analytical HPLC analysis

Method A; A Waters XTerra MS C18 analytical column (3.5 μm , 3.0 \times 150 mm) was used with a flow rate of 0.5 mL/min. A linear gradient of water (containing 0.05% acetic acid) and methanol (containing 0.05% acetic acid) was used, going from 10% organic solvent to 70% organic solvent over a period of 60 min.

Method B; A Waters Xbridge C18 analytical column (3.5 μm , 4.6 \times 150 mm) was used with a flow rate of 0.8 mL/min. A linear gradient of 10 mM ammonium acetate (pH = 5.5) and methanol was used, going from 40% to 60% organic solvent over a period of 45 min.

Purification of compounds with preparative HPLC

Ligands **2-4**, **7** and **8** were purified on a Waters XTerra MS C18 preparative column (3.5 μm , 4.6 \times 150 mm) was used with a flow rate of 0.5 mL/min. A linear gradient of water (containing 0.05% acetic acid) and methanol (containing 0.05% acetic acid) was used, going from 10% organic solvent to 70% organic solvent over a period of 60 min.

Ligands **5** and **6** were purified on a Waters Xbridge C18 preparative column (3.5 μm , 4.6 \times 150 mm) was used for this purpose at a flow rate of 10 mL/min. A linear gradient of 10 mM ammonium acetate (pH = 5.5) and methanol was used, going from 40% to 60% organic solvent over a period of 45 min. The fractions were analysed using analytical HPLC with method A or B. After pooling the fractions containing pure material, the resulting solution was concentrated, extracted with CH_2Cl_2 and dried over Na_2SO_4 . Filtration and evaporation of the solvent yielded pure material.

1-[(2-[(2,5-dioxo-1-pyrrolidinyl)oxy]carbonyl)benzoyl]oxy]-2,5-pyrrolidinedione (10**)**

To a cooled (0°C) solution of *N*-hydroxysuccinimide (1.27 g; 11.0 mmol) in THF (30 mL) was added triethyl amine (1.11 g; 1.52 mL; 11.0 mmol) and freshly distilled *o*-phthaloyl dichloride (1.02 g; 0.73 mL; 5.00 mmol). The resulting white suspension was stirred for 2 hours at room temperature. The solvent was removed and the residue was taken up in dichloromethane (100 mL), washed with water (3 \times 50 mL) and dried over Na_2SO_4 . Filtration and evaporation of the solvent yielded an off-white solid, which was recrystallized from isopropyl alcohol to yield a white solid (1.40 g; 3.90 mmol; 78%). M.p. 161.9-162.0°C.

^1H NMR (300 MHz, CDCl_3) 8.01-8.06 (m, 2H), 7.71-7.78 (m, 2H), 2.85 (s, 8H).

^{13}C NMR (50.3 MHz, CDCl_3) 168.7, 161.5, 133.2, 130.9, 127.3, 25.7.

MS-Cl^+ : m/z 378.1 $[\text{M} + \text{NH}_4]^+$ (100%).

Anal. calcd (%) for $\text{C}_{16}\text{H}_{12}\text{N}_2\text{O}_8$: C 53.34, H 3.36, N 7.78; found: C 53.0, H 3.34, N 7.68.

1-[(3-[(2,5-dioxo-1-pyrrolidinyl)oxy]carbonyl)benzoyl]oxy]-2,5-pyrrolidinedione (11**)**

Following the procedure as for **10** starting from isophthalic acid (2.00 g; 12.0 mmol), DCC (5.52 g; 26.5 mmol) and *N*-hydroxysuccinimide (3.05 g; 26.5 mmol) in THF (300 mL). A white solid (3.78 g; 10.5 mmol; 86%) was obtained after recrystallization from isopropyl alcohol. M.p. 255°C (decomp.).

¹H NMR (400 MHz, CDCl₃) 8.90 (t, *J* = 1.4 Hz, 1H), 8.44 (dd, *J* = 7.9 Hz, *J* = 1.4 Hz, 2H), 7.72 (t, *J* = 7.9 Hz, 1H), 2.93 (s, 8H).

¹³C NMR (50.3 MHz, CDCl₃) 168.8, 160.6, 136.3, 132.4, 129.7, 126.3, 25.6.

MS-Cl⁺: *m/z* 378.1 [M + NH₄⁺]⁺.

Anal. calcd (%) for C₁₆H₁₂N₂O₈: C 53.34, H 3.36, N 7.78; found: C 53.5, H 3.35, N 7.75.

1-[(4-[(2,5-dioxo-1-pyrrolidinyl)oxy]carbonyl]benzoyl)oxy]-2,5-pyrrolidinedione (12**)**

Following the procedure as for **10**, starting from terephthalic acid (2.01 g; 12.1 mmol), DCC (5.49 g; 26.6 mmol) and *N*-hydroxysuccinimide (3.06 g; 26.6 mmol) in THF (300 mL). A white solid (3.74 g; 10.4 mmol; 86%) was obtained after recrystallization from isopropyl alcohol. M.p. 260°C (decomp.).

¹H NMR (400 MHz, CDCl₃) 8.26 (s, 4H), 2.92 (s, 8H).

¹³C NMR (100 MHz, CDCl₃) 168.75, 160.83, 130.81, 130.66, 25.68.

MS-Cl⁺: *m/z* 378.1 [M + NH₄⁺]⁺ (100%).

Anal. calcd (%) for C₁₆H₁₂N₂O₈: C 53.34, H 3.36, N 7.78; found: C 53.4, H 3.44, N 7.73.

1-[(3,5-bis{[(2,5-dioxo-1-pyrrolidinyl)oxy]carbonyl}benzoyl)-2,5-pyrrolidinedione (13**)**

A mixture of 1,3,5-benzene tricarboxylic acid (1.01 g; 4.79 mmol), *N*-hydroxysuccinimide (1.82 g; 15.8 mmol) and DCC (3.30 g; 16.01 mmol) in THF (250 mL) was stirred overnight. The white suspension was filtered and the supernatant was concentrated. The resulting white solid was recrystallized from isopropanol. The product was obtained as a white solid (1.82 g; 3.64 mmol; 76%). M.p. 255°C (decomp.).

¹H NMR (400 MHz, CDCl₃) 9.14 (s, 3H), 2.94 (s, 12H).

¹³C NMR (100 MHz, CDCl₃) 168.50, 159.54, 137.36, 127.67, 25.61.

MS-Cl⁺: *m/z* 388 [M – NHS + H⁺]⁺.

Anal. calcd (%) for C₂₁H₁₅N₃O₁₂: C 50.3; H 3.02, N 8.38; found: C 50.1; H 3.12; N 8.28.

6-benzyl 1-ethyl 3-[3(benzyloxy)-3-oxopropyl]-2-hexenedioate (15**)**

Triethyl phosphonoacetate (3.1 g; 13.84 mmol) in THF (10 mL) was added slowly to a cooled solution of NaH (332 mg; 13.84 mmol) in THF (20 mL). The resulting reaction mixture was stirred for an additional 30 min at 0°C, before benzyl protected 4-ketopimelic acid (**14**) (4.46 g; 12.58 mmol) was added. The resulting mixture was stirred for 1 h at room temperature and heated under reflux overnight. Water (20 mL) was added and the mixture was extracted with EtOAc (3× 20 mL). The organic layers were dried (Na₂SO₄), filtered and the solvent was evaporated. The product was further purified by column chromatography (SiO₂; EtOAc: Pentane; 1: 10) to afford the product **15** (2.89 g; 6.82 mmol; 54%) as a white solid.

¹H NMR (400 MHz, CHCl₃) 7.36 (m, 10H), 5.63 (s, 1H), 5.09 (s, 4H), 4.11 (q, *J* = 7.0 Hz, 2H), 2.87 (t, *J* = 6.2 Hz, 4H), 2.50 (t, *J* = 6.2 Hz, 4H), 1.24 (t, *J* = 7.0 Hz, 3H).

¹³C NMR (50.4 MHz, CDCl₃): δ 206.8, 206.6, 172.5, 172.3, 135.7, 128.4, 128.1, 128.0, 66.3, 60.5, 36.9, 36.9, 31.8, 29.6, 29.2, 27.8, 27.6, 22.5, 14.0.

MS-Cl⁺: *m/z* 442 [M + NH₄⁺]⁺ (100%), 425 [M + H⁺]⁺ (100%).

4-(2-ethoxy-2-oxoethyl)heptanedioic acid (16**)**

To a solution of **15** (1.00 g, 2.36 mmol) in MeOH (25 mL) was added Pd/C (10%, 0.1 g) and the suspension was stirred under a H₂ atmosphere overnight. The mixture was filtered over Celite®. After washing the filtercake with MeOH, the solvent was evaporated to afford the reduced diacid **16** (0.54 g; 2.34 mmol; 99%) as a solid.

¹H NMR (400 MHz, CHCl₃) 4.12 (q, *J* = 7.1 Hz, 2H), 2.23-2.47 (m, 6H), 1.90 (m, 1H), 1.66 (m, 4H), 1.23 (t, *J* = 7.0 Hz, 3H)

¹³C NMR (50.3 MHz, CDCl₃) 177.15, 170.54, 59.51, 38.35, 33.98, 31.21, 28.56,

HRMS (EI⁺) calcd. for C₁₁H₁₇O₅ (M – OH)⁺: *m/z* 229.107; found: 229.106.

4-(carboxymethyl)heptanedioic acid (17**)**

A mixture of **16** (0.21 g; 0.85 mmol) and LiOH (61.33 mg; 2.56 mmol) in MeOH/H₂O (5 mL; 1:1 v/v) was stirred overnight. The mixture was washed with CH₂Cl₂ (10 mL) and conc. H₂SO₄ was added until pH ~ 2. The solution was extracted with EtOAc (12× 10 mL). After drying (Na₂SO₄) and filtration the product was isolated as a white solid (105 mg; 0.48 mmol; 56%).

¹H NMR (300 MHz, CD₃OD) 2.33 (t, *J* = 7.80, Hz, 4H), 2.27 (d, *J* = 6.63 Hz, 5H), 1.89 (td, *J* = 12.94, 6.47 Hz, 1H), 1.65 (m, 4H)

¹³C NMR (100 MHz, CDCl₃) 205.4, 204.6, 67.1, 63.3, 60.1, 57.7.

HRMS (EI⁺) calcd. for C₉H₁₃O₅ (M – OH)⁺: *m/z* 201.076; found: 201.076.

Dimethyl 5-{3-[(*tert*-butoxycarbonyl)amino]propoxy}isophthalate (21**)**

Dimethyl 5-hydroxyisophthalate (1.00 g; 4.73 mmol) was dissolved in acetone (50 mL). To this solution the Boc-protected bromopropylamine **20** (0.563 g; 2.36 mmol) and K₂CO₃ (0.980 g; 7.09 mmol) were added and the white suspension was heated under reflux overnight. After cooling to room temperature the suspension was filtered and the solvent was removed. CH₂Cl₂

(50 mL) was added and after an additional filtration, the solution was washed with sat. aq. NaHCO₃ (2 × 50 mL) and water (50 mL). Drying (Na₂SO₄), filtration and evaporation yielded a colorless oil. After purification by column chromatography (SiO₂; pentane: EtOAc 4:1 to 1:1; R_f = 0.25) a white solid was obtained (0.505 g; 1.37 mmol; 58%). M.p. 76.8–77.0°C.

¹H NMR (200 MHz, CDCl₃) 8.26 (t, J = 1.4 Hz, 1H), 7.72 (d, J = 1.4 Hz, 2H), 4.88–4.58 (br. s, 1H), 4.09 (t, J = 6.0 Hz, 2H), 3.91 (s, 6H), 3.33 (m, 2H), 2.01 (m, 2H), 1.43 (s, 9H).

¹³C NMR (50.3 MHz, CDCl₃) 166.0, 158.8, 155.9, 131.7, 123.0, 119.7, 79.3, 66.2, 52.3, 37.8, 29.4, 28.3.

HRMS (EI⁺) calcd. for C₁₈H₂₅NO₇: *m/z* 367.163; found 367.164

Anal. calcd (%) for C₁₈H₂₅NO₇: C 58.84, H 6.86, N 3.81; found: C 59.0, H 6.78, N 3.61.

5-{3-[(*tert*-butoxycarbonyl)amino]propoxy}isophthalic acid (**18**)

To a solution of the diester **21** (0.525 g; 1.43 mmol) in THF/H₂O (20 mL; 1:1 v/v) was added LiOH (0.103 g; 4.29 mmol). The mixture was stirred at room temperature for 48 h. Conc. H₂SO₄ (96%; 0.24 mL; 4.3 mmol) was added, resulting in a white suspension, which became clear after standing overnight (pH < 1). The solution was extracted with CH₂Cl₂ (4 × 10 mL). The combined organic fractions were washed with water (3 × 10 mL) and dried over Na₂SO₄. Filtration and evaporation yielded a white solid (0.404 g; 1.19 mmol; 83%). M.p. 270.6–272.4°C (decomp.).

¹H NMR (400 MHz, DMSO-*d*₆) 8.04 (t, J = 1.4 Hz, 1H), 7.60 (d, J = 1.4 Hz, 2H), 6.88 (t, J = 5.2 Hz, 1H), 4.05 (t, J = 6.2 Hz, 2H), 3.07 (m, 2H), 1.81 (m, 2H), 1.34 (s, 9H).

¹³C NMR (50.3 MHz, DMSO-*d*₆) 166.4, 158.8, 155.6, 132.6, 122.1, 119.0, 77.5, 65.9, 36.7, 29.1, 28.2.

HRMS (EI⁺) calcd. for C₁₆H₂₁NO₇: *m/z* 339.132; found: 339.132

Anal. calcd (%) for C₁₆H₂₁NO₇: C 56.6, H 6.24, N 4.13; found: C 56.7, H 6.25, N 4.15.

Dimethyl-2-methoxyisophthalate (**22**)

To a solution of 2-methoxyisophthalic acid (5.07 g; 25.8 mmol) in 250 mL MeOH was added 1 mL H₂SO₄. The solution was heated under reflux overnight. After cooling to room temperature the solvents were removed and 50 mL CH₂Cl₂ was added. The solution was washed with H₂O (2 × 50 mL) and dried over Na₂SO₄. Evaporation of the solvent and purification by column chromatography (SiO₂, pentane: EtOAc 9:1 → 5:1 → 3:1) gave **1** (4.37 g, 19.5 mmol, 75%) as a yellow oil.

¹H-NMR (400 MHz, CDCl₃) 7.94 (d, J = 7.7, 2H), 7.21 (t, J = 7.7, 1H), 3.93 (s, 9H).

¹³C NMR (50.3 MHz, CDCl₃) 166.0, 159.5, 134.9, 126.5, 123.4, 63.6, 52.3.

HRMS calcd. for C₁₁H₁₂O₅: *m/z* 224.0684, found: 224.0671.

Dimethyl-2-hydroxyisophthalate (**23**)

A solution of **22** (2.60 g; 11.6 mmol) in 50 mL dry CH₂Cl₂ was placed under N₂-atmosphere and cooled to -78°C. Slowly 1 M BBr₃ in CH₂Cl₂ (12.7 mL; 12.7 mmol) was added. After stirring for 45 minutes at -78°C, 15 mL MeOH was added and the solution was warmed up to room temperature. The mixture was washed with sat. aq. NaHCO₃ (3 × 30 mL) and dried over Na₂SO₄. Evaporation of the solvent gave **23** (1.57 g; 64%) as a yellow oil.

¹H-NMR (CDCl₃) 11.81 (s, 1H), 8.03 (d, J = 7.9, 2H), 6.92 (t, J = 7.9, 1H), 3.93 (s, 6H).

¹³C-NMR (50.3 MHz, CDCl₃) 168.0, 161.5, 136.3, 118.4, 116.4, 52.4.

HRMS calcd. for C₁₀H₁₀O₅: 210.0528; found: 210.0523.

Anal. calcd. C₁₀H₁₀O₅: C, 57.14, H, 4.80; found C, 56.91, H, 4.79.

Dimethyl-2-(3-(*tert*-butoxycarbonylamino)propoxy)isophthalate (**24**)

To a solution of **23** (1.1 g; 4.7 mmol) and K₂CO₃ (3.24 g; 23.5 mmol) in 150 mL DMF was added a solution of **20** (1.0 g; 4.7 mmol) in 10 mL DMF. After heating overnight, the solution was decanted. The solvent was removed and 50 mL CH₂Cl₂ was added. The mixture was washed with brine (3 × 20 mL) and H₂O (3 × 20 mL) and dried over Na₂SO₄. Evaporation of the solvent and purification by column chromatography (SiO₂; pentane: EtOAc 5:1 → 3:1) gave **24** as a yellow oil (1.30 g; 3.5 mmol; 75%).

¹H-NMR (400 MHz, CDCl₃) 7.95 (d, J = 7.8, 2H), 7.18 (d, J = 7.8, 1H), 5.49 (s, 1H), 4.07 (t, J = 5.6, 2H), 3.92 (s, 6H), 3.38 (q, J = 5.9, 11.9 Hz, 2H), 1.96 (m, 2H), 1.45 (s, 9H).

¹³C-NMR (50.3 MHz, CDCl₃) 166.1, 158.8, 156.4, 135.4, 126.8, 123.8, 78.9, 74.2, 52.8, 37.6, 30.1, 28.7.

MS (CI⁺, NH₃) *m/z* 385 ([M + NH₃]⁺, 100), 368 (71), 329 (49), 268 (19).

2-(3-(*tert*-butoxycarbonylamino)propoxy)isophthalic acid (**19**)

To a solution of **23** (0.92 g; 2.37 mmol) in 20 mL THF and 20 mL H₂O was added LiOH (212 mg; 710 mmol). After stirring at room temperature for 24 hours, conc. H₂SO₄ (3.5 mL; 710 mmol) was added. The solution was extracted with CH₂Cl₂ (5 × 20 mL), washed with H₂O and dried over Na₂SO₄. Evaporation of the solvent gave **19** as a white solid (300 mg; 0.88 mmol, 68%). M.p. 148–149°C.

¹H-NMR (400 MHz, DMSO) 7.80 (m, 2H), 7.25 (m, 1H), 6.75 (s, 1H), 3.95 (t, J = 6.0 Hz, 2H), 3.09 (q, J = 6.6, 13.4 Hz, 2H), 1.78 (m, 2H), 1.37 (s, 9H).

¹³C-NMR (50.3 MHz, DMSO) 167.8, 157.0, 156.2, 134.1, 128.6, 124.3, 78.2, 74.3, 38.9, 31.0, 28.9.

MS-CI⁺: *m/z* 357 [M + NH₄]⁺, 340 [M + H]⁺.

Anal. calcd. (%) for C₁₆H₂₁NO₇: C 56.6, H 6.24, N 4.13; found: C 57.1, H 6.47, N 4.02.




Figure S1. ^1H -NMR titration of **3** in D_2O with $(\text{NH}_3)_2\text{Fe}^{\text{II}}(\text{SO}_4)_2$ (—) 0 eq. (—) 1 eq. (—) 2 eq. (—) 3 eq. of $(\text{NH}_3)_2\text{Fe}^{\text{II}}(\text{SO}_4)_2$. Inset; plot of the relative area of the peak at δ 25.9 ppm compared to the acetone peak against the amount of equivalents of $(\text{NH}_3)_2\text{Fe}^{\text{II}}(\text{SO}_4)_2$.




Figure S2. UV/Vis spectra of ligand **3** after addition of (—) 1 eq. (—) 2 eq. (—) 3 eq. of $(\text{NH}_3)_2\text{Fe}^{\text{II}}(\text{SO}_4)_2$. Inset; plot of the $\epsilon/\epsilon_{\text{max}}$ at 356 nm against the amount of equivalents of $(\text{NH}_3)_2\text{Fe}^{\text{II}}(\text{SO}_4)_2$.




Figure S3 Aerobic oxidation of supercoiled plasmid DNA (■) into nicked DNA (●) and linear DNA (▲) followed in time catalyzed by a) $[(2)\text{Fe}^{\text{II}}_2]^{4+}$ (**30**) and b) $[(4)\text{Fe}^{\text{II}}_2]^{4+}$ (**32**). Errors bars represent the root mean square (rms) error based on three runs. A correction factor of 1.31 was used to compensate for the reduced ethidium bromide uptake capacity of supercoiled DNA







Figure S4: Example of an agarose gel slab of the reaction of supercoiled DNA to form nicked and linear DNA with complex **32**. Each lane represents a surtain time point in the time profile plots; lane 1, 20s; lane 2, 40s; lane 3, 1 min.; lane 4, 2 min.; lane 5, 3 min.; lane 6, 4 min.; lane 7, 5 min.; lane 8, 7.5 min.; lane 9, 10 min.; lane 10, 15 min.; lane 11, 20 min.; lane 12, 25 min.; lane 13, 30 min.; lane 14, 45 min; lane 15, 60 min.


¹H-NMR




¹³C-NMR




Analytical RP-HPLC




¹H-NMR




¹³C-NMR




Analytical RP-HPLC




¹H-NMR




¹³C-NMR




Analytical RP-HPLC




$^1\text{H-NMR}$




$^{13}\text{C-NMR}$




Analytical RP-HPLC




¹H-NMR




¹³C-NMR




Analytical RP-HPLC



¹H-NMR



¹³C-NMR



Analytical RP-HPLC

